Potential use of corn co-products in fishmeal-free diets for juvenile Nile tilapia *Oreochromis niloticus*

著者	Sandamali Sakunthala Herath, 芳賀 穰, 佐藤 秀一
journal or publication title	Fisheries Science
volume	82
number	5
page range	811-818
year	2016-09
権利	(c) 2016 Japanese Society of Fisheries Science and Springer Japan. This is the author's version of the work. It is posted here for your personal use. Not for redistribution. The definitive Version of Record was published in https://doi.org/10.1007/s12562-016-1008-6
科学研究費研究課題	魚類の育成環境適応性と環境負荷低減を考慮した養魚飼料の基礎的研究 Studies on the development of aquafeed with consideration of rearing condition and low load to environment
研究課題番号	26252034
URL	http://id.nii.ac.jp/1342/00001472/

doi: 10.1007/s12562-016-1008-6

[Click here to view linked References](#)

Fisheries Science 82(5) 811-818, doi

1
2
3
4
5
6 1 **Potential use of corn co-products for fishmeal-free diets for juvenile Nile tilapia *Oreochromis niloticus***

7
8 2 Sandamali Sakunthala Herath^{1,2}, Yutaka Haga¹ & Shuichi Satoh¹

9
10
11 3 ¹ Graduate School of Marine Science and Technology, Tokyo University of Marine Science and Technology,

12
13 4 Tokyo, Japan

14
15
16 5 ² Department of Fisheries & Aquaculture, Faculty of Fisheries and Marine Sciences & Technology,

17
18 6 University of Ruhuna, Matara, Sri Lanka

19
20
21 7

22
23
24 8 Correspondence: Shuichi Satoh

25
26
27 9 Tel: +81-3-5463-0557

28
29
30 10 Fax: +81-3-5463-0553

31
32
33 11 E-mail addresses: sakunthala@fish.ruh.ac.lk, haga@kaiyodai.ac.jp, ssatoh@kaiyodai.ac.jp

34
35
36 12

37
38
39 13

40
41
42 14

43
44
45 15

46
47
48 16

49
50
51 17

52
53
54 18

55
56
57 19

58
59
60 20

- 1 ティラピア *Oreochromis niloticus* 稚魚用無魚粉飼料におけるトウモロコシ副産物
2 の有効性
- 3 Sandamali Sakunthala Herath (Ruhuna 大学, 海洋大) , 芳賀穰, 佐藤秀一 (海洋大)
- 4 ティラピア *Oreochromis niloticus* 稚魚用無魚粉飼料におけるトウモロコシ副産物の
5 有効性を評価した。すなわち、魚粉または濃縮トウモロコシタンパク、コーング
6 ルテンミール、高タンパクトウモロコシ蒸留粕 (HPDDG) 、可溶性物含有トウモ
7 ロコシ蒸留粕(DDGS)を含む飼料を作製し、4.5 g の魚に 12 週間給餌した。対照区
8 および DDGS 飼料を給餌区で有意に高い飼育成績が得られ、次いで HPDDG 区で
9 優れた成績が得られた。また、全魚体と筋肉中タンパク質含量も HPDDG 区で最
10 も高く、脂質含量は DDGS 区で高かった。DDGS はティラピア用飼料の魚粉を完
11 全に代替するのに有効であると示唆された。

21 **Abstract**

22 We conducted a 12-week feeding trial to evaluate the effects of total fishmeal replacement with
23 different corn co-products on growth performance, feed utilization efficiency and body composition in
24 juvenile Nile tilapia *Oreochromis niloticus*. Five isonitrogenous diets were prepared. Fifty per cent of the
25 dietary protein was obtained from fishmeal (control diet) or from one of four corn co-products, namely corn
26 protein concentrate, corn gluten meal, high-protein distillers' dried grains (HPDDG) or distillers' dried grains
27 with solubles (DDGS) (experimental diets). Fish with an initial mean weight of 4.5 g were fed one of the five
28 diets twice a day to near satiety. Significantly higher ($P < 0.05$) specific growth rates and survival occurred in
29 fish fed the control diet or DDGS, followed by HPDDG; mean feed intakes by fish in these three groups were
30 significantly greater than those in the others. Food conversion ratio, protein efficiency ratio, and total amino
31 acid content of whole body were not affected by changes in dietary ingredients. Whole body and fillet protein
32 content was highest in the HPDDG group, and lipid content was highest in those fed DDGS. DDGS can be
33 used to fully replace the fishmeal component of Nile tilapia diets.

35 Key words: amino acids, DDGS, feed utilization, fillet quality, growth performance, HPDDG

1. Introduction

The search for alternatives to replace the fishmeal component of aqua-feed has received a good deal of attention, resulting in considerable research progress, over the last two decades. A vast array of proteins from both plant and animal sources has been evaluated widely for their suitability for partial or total replacement of fishmeal in aqua-feeds [1-7]. Among the tested ingredients, industrial by-products or co-products that are considered unsuitable for direct human consumption have received much interest [8]. In this context, protein-rich co-products of the corn-milling industry play an important role as protein sources for the manufacturing of animal feeds, including aqua-feeds.

Corn gluten meal (CGM) and corn protein concentrate (CPC) are co-products of the corn wet-milling industry, and CGM has been used widely in aqua-feeds[9]. Distillers' dried grains with solubles (DDGS) and high-protein distillers' dried grains (HPDDG) are co-products of corn dry-milling industries for fuel ethanol production. HPDDG is a relatively new product of the corn ethanol industry, and its nutritional value is much more consistent than that of DDGS [10] . Unlike other, conventional, plant protein sources such as soybean meal and cotton seed meal, corn co-products are free from anti-nutritional factors [11, 12]and few amino acids deficiencies were reported [13]. However, differences among variety of corn co-products, such as percentage protein can be identified due to the differences in the wet-milling and dry-milling processes.

Aquaculture of tilapia, a group of fish with herbivorous or omnivorous feeding habits is the most widespread in the world. Dietary substitution of fishmeal with an alternative protein source in herbivorous or omnivorous species is considerably easier than in carnivorous species, which are nutritionally more demanding[14].

Because tilapia can utilize a high percentage of dietary plant ingredients [15], co-products of the corn-milling industry such as CGM and DDGS have been tested in their diets, with varying degree of success [1-3, 5, 16, 17]. However, total fishmeal replacement with a single corn co-product has not yet been evaluated in this fish, and to our knowledge no attempt has been made to compare multiple corn co-products

as fishmeal alternatives in a single growth trial. Our objectives here were to 1) compare the effects of total fishmeal replacement with various single corn co-products in the diet of Nile tilapia *Oreochromis niloticus*, juveniles, on growth performance, feed utilization efficiency, and body composition; and 2) determine the most suitable corn co-product for use in a zero-fishmeal diet for these fish.

2. Materials and Methods

2.1 Experimental diets

Five isonitrogenous (32% protein) experimental diets (fishmeal-based control diet and four corn co-product-based diets) were formulated to fulfil the known nutritional requirements of juvenile Nile tilapia [18]. Proximate composition of main protein sources and ingredient composition of diets are given in Table 1 and 2 respectively. In the control diet, 50% of the protein was supplied by fishmeal. In the other four diets, the fishmeal component of the control diet was fully replaced with one of four corn co-products, namely CPC, CGM, HPDDG or DDGS. Crystalline amino acids (DL- methionine & L-lysine) were used to compensate the amino acid deficiencies in corn based diets.

< Table 1-2 >

The proximate composition and total amino acid composition of the experimental diets is presented in Table 3. Before feed preparation, the ingredients were ground to pass through a mesh (0.5 µm) screen and then mixed by using a horizontal mixer. The mixture was then moistened by adding distilled water (~20%), pelleted and freeze-dried. Diets were stored at 5 °C until use. The names of the five diets were designated according to the main protein ingredient, namely control (FM), CPC, CGM, HPDDG and DDGS.

<Table 3 >

2.2 Experimental fish and rearing condition

Nile tilapia fingerlings were obtained from the Laboratory of Fish Culture, Tokyo University of Marine Science and Technology, Tokyo, Japan. Before the start of the experiment, the fish were acclimated to the experimental conditions for two weeks. During this acclimation period, fish were hand fed to near satiety twice a day by using the control diet. After the acclimatization period food was withheld for 24 h, after which 200 fish with similar initial body weights (4.0 to 4.5 g) were selected for potential use in the experiment. Groups of 18 fish were bulk weighed and stocked into each glass tank (60 L). Five treatments, one for each diet (control, CPC, CGM, HPDDG and DDGS), in duplicate were randomly allocated to ten (2 × 5) glass tanks. Each tank was connected to a freshwater re-circulating system in which the water temperature was maintained at 28 ± 1 °C by using electrical heaters (immersion heaters). The water flow rate was maintained at 0.5 L min⁻¹, and aeration was provided continuously to each tank via submerged air stones. Approximately 50% of the water in the re-circulating system was replaced with de-chlorinated tap water once every two weeks. Fish were hand fed to near satiety twice a day, 6 days a week for 12 weeks, and daily feed intakes were recorded.

2.3 Data collection and sampling

The 20 fish remaining in the initial stock of 200 were sacrificed by using an overdose of 2-phenoxyethanol and kept at -30 °C for initial carcass analysis. During the 12-week experiment, the fish in each tank were bulk weighed at 3-week intervals. At the end of the experiment, fish were starved for 24 h and anaesthetized with 0.2% 2-phenoxyethanol before being handled. The weight and length of the fish in each tank were individually measured with an electrical balance and a measuring board, respectively, to the nearest first decimal point.

Five average-size fish from each tank were euthanized using overdose of 2-phenoxyethanol and used to calculate the hepatosomatic index (HSI), viscerosomatic index (VSI) and fillet yield (FY). Four fish from each treatment and the fillet samples used for fillet yield determination were immediately frozen at -30 °C for final chemical analysis. Frozen samples were minced, freeze-dried and kept at -30 °C until analysis.

Percentage specific growth rate (SGR, %), thermal growth coefficient (TGC), food conversion ratio (FCR), protein efficiency ratio (PER), Protein retention (%), survival rate (SR, %), HSI, VSI, fillet yield and the coefficient of condition (K) were calculated by using the following equations to compare fish growth, nutrient utilization efficiency, and body indices among treatments.

$$SGR = (\ln FW - \ln IW) / \text{no. of days} \times 100$$

$$TGC = [(FW^{1/3} - IW^{1/3}) / (\text{water temperature } ^\circ\text{C} \times \text{no. of days})] \times 1000$$

$$FCR = \text{feed intake (dry basis)} / \text{wet weight gain}$$

$$PER = \text{body weight gain (g)} / \text{protein intake (g)}$$

$$\text{Protein retention (\%)} = (\text{final body protein} - \text{initial body protein}) / \text{protein intake} \times 100$$

$$SR = \text{number of fish at harvest} / \text{number of fish stocked} \times 100$$

$$HSI = \text{liver weight} / \text{body weight} \times 100$$

$$VSI = \text{visceral weight} / \text{body weight} \times 100$$

$$\text{Fillet yield} = \text{fillet weight} / \text{body weight} \times 100$$

$$K = 100 \times \text{mean weight (g)} / (\text{total length (cm)})^3$$

FW, final mean weight of fish; IW, initial mean weight of fish

2.4 Chemical analysis

Feed ingredients, diets, whole body and muscle samples of fish were analysed in accordance with standard procedures for chemical analysis [19]. All the samples were finely ground and analysed in triplicate. Dry matter content was calculated from the weight loss after drying of the sample at 105 °C until it reached a

constant weight. Ash content was determined after the incineration of samples in a muffle furnace at 550 °C for 16 h. The Kjeldhal method was used for crude protein analysis. Crude lipid content was determined by the gravimetric method after extraction of the lipids into a mixture of chloroform and methanol (2:1 v/v).

Amino acid content was analysed by using an automatic amino acid analyser (JLC-500/v; JEOL, Tokyo, Japan)[20]. For total amino acids, samples were digested at 110 °C for 22 h with 4 M methanosulphonic acid (Sigma-Aldrich, St. Louis, MO, USA); the digested solution was then passed through a 0.45-µm membrane filter and injected into the analyser.

2.5 Statistical analysis

Data were initially checked for normality and equal variance by using Levene's test for equality of variances. The effects of different corn co-products on Nile tilapia growth performance, feed utilization efficiency, body indices and whole body and fillet proximate composition were compared in a one-way analysis of variance (ANOVA) at the 5% significance level ($P < 0.05$). When ANOVA was found to be significant, Tukey's multiple range tests was used to detect differences among treatments. All statistical analyses were performed with SPSS version 16.0.

3. Results

3.1 Growth performances and feed utilization efficiencies

The initial weights of fish assigned to the different treatments (4.0 to 4.5 g) did not differ significantly ($P > 0.05$). However, final weight was significantly affected by the treatments (Fig. 1).

<Fig. 1>

Differences in the growth rates of fish fed various diets were first observed after 6 weeks. The final mean weights of fish in the five treatment groups differed significantly from each other ($P < 0.05$). The

greatest final weight was observed in the control, followed by DDGS. The lowest weight was observed in the CPC treatment. Among the corn-based test diets, DDGS gave a significantly greater final mean weight, followed by HPDDG. Final mean weight of DDGS was almost double than that of CPC. The weights of fish in the CPC and CGM groups almost overlapped until week 9, but by the end of the experiment CGM had resulted in a significantly greater weight than did CPC. Growth performance and feed utilization efficiencies of the fish are given in Table 4.

<Table 4>

The % SGRs of the control and DDGS groups were significantly higher than those of the others. The lowest SGR was observed in the CPC group, followed by the CGM group. TGC differed significantly among treatments and it was highest in the control. Mean feed intake was significantly higher in the control, DDGS and HPDDG groups than in the others. Feed utilization efficiency in terms of FCR and protein efficiency ratio (PER) was independent of the type of dietary treatment. However, protein retention was significantly affected by the treatment. Protein retention in control was significantly higher than that of CPC and CGM while HPDDG and DDGS differ only from CPC. Survival rate was significantly affected by the treatment. The CPC, CGM and HPDDG groups had similar survival rates; they were significantly lower than that of the control. However, the survival rate of the DDGS group did not differ from those of the control or HPDDG group.

3.2 Whole body and fillet proximate composition

We examined the proximate compositions of the whole body and fillet (Table 5). The dry matter content of the whole body was not influenced by diet. Crude protein content of fish whole body was significantly higher in HPDDG than in the other groups, with the next-highest contents in the control and DDGS groups; these three groups had significantly higher feed intakes than the other treatment groups (see Table 4), and the control and DDGS groups had significantly higher growth rates than the others. Whole body lipid content in all the corn-based treatments was significantly higher than that of the control. Ash

content was also greatly affected by dietary treatment; it was significantly higher in the control than in the other groups; the next-highest value was in the DDGS group.

<Table 5>

Our data on the proximate analysis of fillet samples revealed no treatment effects on dry matter and ash. In the case of both whole body and fillet, the crude protein content was significantly higher in the HPDDG group than in the others, whereas the fillet crude lipid content was highest in DDGS. However, in the whole body or fillet the amount of protein in the control did not differ from that in DDGS; these two groups also had similar growth performances. Similar to the case in the whole body, the lowest crude lipid content in the fillet was observed in the fishmeal-based control.

We examined the total amino acid compositions of whole body at the end of the 12-week period (Table 6); these compositions were independent of treatment type.

<Table 6>

We also examined the morphometric indices of different treatments (Table 7). There were no significant between-group differences in any of the indices evaluated, i.e. VSI, HSI, FY, or K value.

<Table 7>

4. Discussion

Growth in terms of the weight of juvenile fish usually follows a sigmoid curve, as we observed here in Nile tilapia. Although fish in all treatments had similar growth patterns, growth rates differed among all treatments. Growth of fish is a function of feed intake and capacity of fish to utilize ingested food [21]. Since the feed intake in this study showed significant differences among treatments, the observed variations in growth rate were likely explained by the factors affected on feed intake.

Dietary availability of essential nutrients such as amino acids is a factor which determine the feed intake of fish[22].Therefore differences in feed intake in this study can be explained by the dietary essential amino acid composition. Even if diets were formulated to fulfil the essential amino acid requirement of Nile tilapia [23] (estimated using values in [24]), analytical data showed that some amino acids such as arginine, histidine and valine are deficient in CPC and CGM diets. Essential amino acid composition of control, HPDDG and DDGS was almost similar except histidine and those were the treatment group which showed higher growth performances and feed intakes compared to CPC and CGM. Feed intake of fish is known to reduce when dietary essential nutrients are deficient[22, 25]. Therefore differences in amino acid compositions in diets are the most plausible responsible factor for observed variation in feed intakes and thereby for differences in growth rate of fish as previously observed for Nile tilapia [26].

In addition, DDGS and HPDDG, which showed similar feed intake to the fishmeal-based control, are co-products of a corn dry-milling process, whereas CPC and CGM are co-products of corn wet-milling process. Because all four corn co-products were made from the same ingredient, the observed differences in feed intake and growth are likely attributable to difference in the corn processing method. Unlike wet-milling, dry-milling processes the whole corn kernel, which is subjected to subsequent saccharification and fermentation before the co-products are processed [26-28]. Therefore, dry-milling co-products such as DDGS and HPDDG usually contain fermentation residues, including yeast cells (*Saccharomyces cerevisiae*) [28-30] and other unidentified nutrients formed during the fermentation–distillation process [28]. Corn wet-milling uses steeping and fractionation technology to separate the corn kernel into protein, starch, oil, and fiber components, and the protein portion is not subjected to fermentation [26, 28]. Therefore, the greater feed intake and growth performance observed in the DDGS and HPDDG groups than in the CPC and CGM groups could be associated with the availability of yeast and other unidentified fermentation residues remaining in the dry-milled co-products. This hypothesis is supported by the work of [31] who reported improved growth performance of Nile tilapia juveniles fed diets supplemented with commercial live yeast (*S. cerevisiae*); they noted that the yeast enhanced appetite and consequently improved feed intake. Improved

feed consumption and growth associated with diets containing DDGS has previously been reported for the same species [32].

The % SGRs of fish in the control, DDGS and HPDDG groups were similar to those reported for Nile tilapia of similar size by [32, 33] but slightly higher than those observed by [34]. Even though the % SGRs were lower in the CPC and CGM treatments than in the other treatments, our values were consistent with the reported values in the literature for Nile tilapia of similar size reared at 28 °C in fresh water[34, 35].

Our feed utilization efficiencies in terms of FCR and PER were independent of the treatment type despite the large differences in growth parameters. These findings agreed with those of a previous study[26, 33].However, dietary ingredients seemed to affect on protein retention of this study and values are almost similar to what was reported in literature [36]. Fish can utilize organic macronutrients such as protein, lipid and carbohydrate as a direct metabolic fuel or store for utilization in later date or deposit in the structural materials which represent as growth [37].Protein retention in this study showed close relationship with weight gain of fish and indicated the differences in nutrient retention are partly responsible for differences of growth. Although it was difficult to distinguish statistical differences of protein retention among corn based treatments, fishmeal based control which had highest final body weight showed the higher protein retention compared to that of CPC and CGM which had lower final weight. Dietary amino acid deficiency was also observed in these two treatments and therefore, it is obvious that, protein retention was affected by the dietary amino acid composition. However difficulties in finding differences in feed utilization parameters among corn based treatment groups which were significant for growth parameters may be due to low statistical power in this study which arises with limited number of replicates used.

The whole body and fillet proximate compositions of our experimental fish were affected by the dietary ingredients. However, whole body and fillet protein content of DDGS and HPDDG was equal to or higher than that in the control. This result suggests that complete replacement of fishmeal in the tilapia diet with corn dry milling co-products does not negatively affect the final nutritional quality of fish products in terms of body protein. Because the total amino acid content of the whole body did not differ significantly among treatments, complete replacement of fishmeal with corn co-products had no negative impact even

from the perspective of essential amino acids. Whole body protein of CPC and CGM is lower than that of control, DDGS and HPDDG and these statistical differences were not exist in the fillet protein content. Therefore, lower percentage of whole body protein observed in CPC and CGM was not due to the differences in converting feed protein into fish muscle protein. Since whole body protein is the average of all structural protein and muscle protein, observed variation in whole body protein among treatments should be due to the treatment effects on structural proteins. This should be partly associated with the amino acid deficiency of CPC and CGM diets. Our fillet yield varied in a narrow range (28% to 32%) and was not affected by treatment. This indicates that Nile tilapia are capable of converting corn co-products and other ingredients into edible muscle mass at rates similar to those on a fishmeal-based diet.

In conclusion, total fishmeal replacement with different corn co-products had different effects on the growth performance and proximate composition of the fish's whole body and fillets. Among the corn co-products tested, DDGS was the best choice for a zero fishmeal diet for Nile tilapia juveniles. Because we focused here only on the juvenile stage of Nile tilapia with duplicate, a long-term growth trial with more replicates and balanced amino acid profiles in diets should be done to confirm the effects of corn co-products on the growth performance and fillet quality of marketable-size fish.

Acknowledgements

The expenses of this study is partly defrayed from Grant-in-Aid for Scientific Research given to Shuichi Satoh from Ministry of Education, Culture, Sports, Science and Technology through Japanese Society for Promotion of Science (26252034) and "Development of Aquaculture Technology for Food Security and Food Safety for the Next Generation" by JST and JICA under SATREPS program.

References

- 280 1 Wu YV, Rosati R, Sessa DJ ,Brown P (1994) Utilization of protein-rich ethanol co-
281 products from corn in tilapia feed. J Am Oil Chem Soc. 71:1041-1043
- 282 2 Wu YV, Rosati RR ,Brown PB (1996) Effect of diets containing various levels of
283 protein and ethanol coproducts from corn on growth of tilapia fry. J Agric Food
284 Chem. 44:1491-1493
- 285 3 Tidwell JH, Coyle SD, VanArnum A, Weibel C ,Harkins S (2000) Growth,
286 Survival, and Body Composition of Cage-Cultured Nile Tilapia *Oreochromis*
287 *niloticus* Fed Pelleted and Unpelleted Distillers Grains with Solubles in Polyculture
288 with Freshwater Prawn *Macrobrachium rosenbergii*. J World Aquacult.
289 Soc.31:627-631
- 290 4 Richter N, Siddhuraju P ,Becker K (2003) Evaluation of nutritional quality of
291 moringa (*Moringa oleifera* Lam.) leaves as an alternative protein source for Nile
292 tilapia (*Oreochromis niloticus* L.). Aquaculture 217:599-611
- 293 5 Schaeffer TW, Brown ML, Rosentrater KA ,Muthukumarappan K (2010)
294 Utilization of diets containing graded levels of ethanol production co - products by
295 Nile tilapia. J Anim Physiol Anim Nutr. 94:e348-e354
- 296 6 Lu F, Haga Y ,Sato S (2015) Effects of replacing fish meal with rendered animal
297 protein and plant protein sources on growth response, biological indices, and amino
298 acid availability for rainbow trout *Oncorhynchus mykiss*. Fish Sci. 81:95-105
- 299 7 Plaipetch P ,Yakupitiyage A (2014) Effect of replacing soybean meal with yeast -
300 fermented canola meal on growth and nutrient retention of Nile tilapia,
301 *Oreochromis niloticus* (Linnaeus 1758). Aquacult Res. 45:1744-1753
- 302 8 Xu Z, Lin X, Lin Q, Yang Y ,Wang Y (2007) Nitrogen, phosphorus, and energy
303 waste outputs of four marine cage-cultured fish fed with trash fish. Aquaculture
304 263:130-141
- 305 9 Pereira T ,Oliva - Teles A (2003) Evaluation of corn gluten meal as a protein
306 source in diets for gilthead sea bream (*Sparus aurata* L.) juveniles. Aquacult Res.
307 34:1111-1117
- 308 10 Robinson PH, Karges K ,Gibson ML (2008) Nutritional evaluation of four co-
309 product feedstuffs from the motor fuel ethanol distillation industry in the
310 Midwestern USA. Anim Feed Sci Technol. 146:345-352
- 311 11 Shiau S-Y, Chuang J-L ,Sun C-L (1987) Inclusion of soybean meal in tilapia
312 (*Oreochromis niloticus* × *O. aureus*) diets at two protein levels. Aquaculture
313 65:251-261

314	12	Robinson EH ,Li MH (2008) Replacement of Soybean Meal in Channel Catfish, <i>Ictalurus punctatus</i> , Diets with Cottonseed Meal and Distiller's Dried Grains with Solubles. J World Aquacult Soc. 39:521-527
315		
316		
317	13	Cheng ZJ ,Hardy RW (2004) Nutritional value of diets containing distiller's dried grain with solubles for rainbow trout, <i>Oncorhynchus mykiss</i> . J Appl Aquaculture 15:101-113
318		
319		
320	14	Tacon AG ,Metian M (2008) Global overview on the use of fish meal and fish oil in industrially compounded aquafeeds: trends and future prospects. Aquaculture 285:146-158
321		
322		
323	15	Twibell RG ,Brown PB (1998) Optimal Dietary Protein Concentration for Hybrid Tilapia <i>Oreochromis niloticus</i> × <i>O. aureus</i> Fed All-Plant Diets. J World Aquacult Soc. 29:9-16
324		
325		
326	16	Wu YV, Rosati RR, Sessa DJ ,Brown PB (1995) Evaluation of corn gluten meal as a protein source in tilapia diets. J Agric Food Chem. 43:1585-1588
327		
328	17	Coyle SD, Mengel GJ, Tidwell JH ,Webster CD (2004) Evaluation of growth, feed utilization, and economics of hybrid tilapia, <i>Oreochromis niloticus</i> × <i>Oreochromis aureus</i> , fed diets containing different protein sources in combination with distillers dried grains with solubles. Aquacult Res. 35:365-370
329		
330		
331		
332	18	Jobling M (2012) National Research Council (NRC): Nutrient requirements of fish and shrimp. Aquacult Intl. 20:601-602
333		
334	19	Intl A (1995) Official methods of analysis. Arlington, Va.: AOAC Intl
335	20	Boonyoung S, Haga Y ,Sato S (2013) Preliminary study on effects of methionine hydroxy analog and taurine supplementation in a soy protein concentrate - based diet on the biological performance and amino acid composition of rainbow trout [<i>Oncorhynchus mykiss</i> (Walbaum)]. Aquacult Res. 44:1339-1347
336		
337		
338		
339	21	Carter C, Houlihan D, Kiessling A, Médale F ,Jobling M (2001) Physiological effects of feeding. Food intake in fish:297-331
340		
341	22	De la Higuera M (2001) Effects of nutritional factors and feed characteristics on feed intake. Food Intake in Fish 827:250-268
342		
343	23	Santiago CB ,Lovell RT (1988) Amino acid requirements for growth of Nile tilapia. J. Nutr. 118:1540-1546
344		
345	24	Prachom N, Haga Y ,Sato S (2013) Impact of dietary high protein distillers dried grains on amino acid utilization, growth response, nutritional health status and
346		

- 347 waste output in juvenile rainbow trout (*Oncorhynchus mykiss*). Aquacult Nutr.
348 19:62-71
- 349 25 Gómez-Requeni P, Mingarro M, Caldach-Giner J, Médale F, Martin S, Houlihan D,
350 Kaushik S ,Pérez-Sánchez J (2004) Protein growth performance, amino acid
351 utilisation and somatotrophic axis responsiveness to fish meal replacement by plant
352 protein sources in gilthead sea bream (*Sparus aurata*). Aquaculture 232:493-510
- 353 26 Wu YV, Rosati RR ,Brown PB (1997) Use of corn-derived ethanol coproducts and
354 synthetic lysine and tryptophan for growth of tilapia (*Oreochromis niloticus*) fry. J
355 Agric Food Chem. 45:2174-2177
- 356 27 Weigel J, Loy D ,Kilmer L (2005) Feed co-products of the dry corn milling process.
357 Iowa State University and Iowa Corn Promotion Board.
- 358 28 Weigel JC, Loy D, Kilmer LH, Association ICG, Association RF ,Association
359 NCG (1997) Feed co-products of the dry corn milling process. Iowa Corn Growers
360 Association
- 361 29 Belyea RL, Rausch KD ,Tumbleson ME (2004) Composition of corn and distillers
362 dried grains with solubles from dry grind ethanol processing. Bioresource
363 Technology 94:293-298
- 364 30 Han J ,Liu K (2010) Changes in composition and amino acid profile during dry
365 grind ethanol processing from corn and estimation of yeast contribution toward
366 DDGS proteins. J Agric Food Chem. 58:3430-3437
- 367 31 Abdel-Tawwab M, Abdel-Rahman AM ,Ismael NEM (2008) Evaluation of
368 commercial live bakers' yeast, *Saccharomyces cerevisiae* as a growth and
369 immunity promoter for Fry Nile tilapia, *Oreochromis niloticus* (L.) challenged in
370 situ with *Aeromonas hydrophila*. Aquaculture 280:185-189
- 371 32 Lim C, Garcia JC, Yildirim-Aksoy M, Klesius PH, Shoemaker CA ,Evans JJ
372 (2007) Growth Response and Resistance to *Streptococcus iniae* of Nile Tilapia,
373 *Oreochromis niloticus*, Fed Diets Containing Distiller's Dried Grains with Solubles.
374 J World Aquacul Soc. 38:231-237
- 375 33 He J-y, Han B, Tian L-x, Yang H-j, Zeng S-l ,Liu Y-j (2014) The sparing effect of
376 cystine on methionine at a constant TSAA level in practical diets of juvenile Nile
377 tilapia *Oreochromis niloticus*. Aquacult Res. 47:2031-2039
- 378 34 Likongwe JS, Stecko TD, Stauffer Jr JR ,Carline RF (1996) Combined effects of
379 water temperature and salinity on growth and feed utilization of juvenile Nile
380 tilapia *Oreochromis niloticus* (Linnaeus). Aquaculture 146:37-46

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

381 35 Fontainhas-Fernandes A, Gomes E, Reis-Henriques MA ,Coimbra J (1999)
382 Replacement of fish meal by plant proteins in the diet of Nile tilapia: digestibility
383 and growth performance. Aquacult Intl. 7:57-67

384 36 Furuya WM, Pezzato LE, Barros MM, Pezzato AC, Furuya VR ,Miranda EC
385 (2004) Use of ideal protein concept for precision formulation of amino acid levels
386 in fish - meal - free diets for juvenile Nile tilapia (*Oreochromis niloticus* L.).
387 Aquacult Res. 35:1110-1116

388 37 Jobling M (2001) Feed composition and analysis. Food Intake in Fish 827:1-24
389
390

391 **Figure captions**

392 **Fig. 1** Mean weight of fish over experimental period (n=2, different letters indicate significant differences, p
393 < 0.05)

Figure

Table 1 Proximate composition (g kg⁻¹) of protein sources used in experimental diets fed Nile tilapia *Oreochromis niloticus* (n=3)

Ingredient	Dry matter	Protein	Lipid	Ash
FM	918	682	100	113
CPC	901	798	95.8	16
CGM	933	658	130	18
HPDDG	904	461	37.7	38
DDGS	896	314	77.1	66
SBM	934	458	21	63
PBM	877	690	106	101

FM, fish meal (anchovy); CPC, corn protein concentrate, Emphyreal 75®, Cargill Corn Milling; CGM, corn gluten meal; HPDDG, high protein distillers dried grains, Dakota Gold ® BFRAC™, SBM, soybean meal; PBM, poultry by product meal

Table 2 Ingredient composition (g kg⁻¹) of experimental diets fed Nile tilapia *Oreochromis niloticus*

Ingredient	Control	CPC	CGM	HPDDG	DDGS
Fish meal	218	0	0	0	0
Corn co-product	0	194	235	332	524
Soybean meal	162	166	168	188	211
Poultry by product meal	51	53	52	52	71
Wheat flour	312	329	302	225	56
Alpha starch	190	176	163	121	86
Vitamin mix	10	10	10	10	10
Mineral mix	16	16	16	16	16
Soy oil	41	44	44	46	18
L-lysine	0	8	7	6	4
DL-methionine	0	4	3	4	4
Crystalline amino acids	0	12	10	10	8

CPC; corn protein concentrate based diet, CGM; corn gluten meal based diet, HPDDG; high protein distillers dried grain based diet, DDGS; distillers dried grains with soluble based diet

Vitamin mixture composition (unit kg⁻¹premix): Vitamin A, 420 000 IU; Vitamin D3, 2 420 000 IU; Vitamin K3, 6050mg; Vitamin E 1000mg; thiamine, 3025 mg; riboflavin, 3630 mg; pyridoxine, 2420 mg; cyanocobalamine, 6 mg; L-ascorbic acid, 368 900mg; nicotinic acid, 24 200mg; D-pantothenic acid, 6050mg; inositol, 121 000mg; d-biotin, 363 mg; folic acid, 908 mg; para-aminobenzoic acid 3025 mg.

Mineral mixture composition (g/kg⁻¹ premix) : Sodium chloride 50; magnesium sulphate 745; iron(III) citrate n-hydrate 125; trace element mix 50; cellulose 30 [the trace element mixture contains(g/kg⁻¹) Zinc sulphateheptahydrate 353; manganese sulphate 162; copper (II) sulphate pentahydrate 31; aluminium chloride hexahydrate 10; cobalt chloride 3; potassium iodate 1; cellulose 440].

Table 3 Proximate and amino acid composition of experimental diets (g kg⁻¹) fed Nile tilapia *Oreochromis niloticus* (n=3)

Components	Control	CPC	CGM	HPDDG	DDGS
Dry matter	968	963	964	961	961
Protein	319	323	321	321	315
Lipid	88.7	97.2	97.4	88.9	98.4
Ash	76	38.7	34.4	45.2	56.5
EAA					
Arginine	14.8	9.1	10.5	14.1	13.9
Histidine	7.5	3.9	4.4	6.0	5.7
Isoleucine	6.8	5.6	6.5	8.5	7.7
Leucine	17.7	19.8	23.2	23.6	22.8
Lysine	15.0	11.8	12.5	16.6	14.6
Methionine	5.3	5.9	6.1	7.7	8.1
Phenylalanine	11.0	9.9	11.7	12.5	11.6
Threonine	10.3	6.8	8.0	11.0	9.9
Tryptopan	1.0	0.7	0.8	1.0	0.6
Valine	8.4	5.7	7.5	10.4	9.5
NEAA					
Alanine	15.4	12.5	14.9	16.1	15.6
Aspartic acid	25.5	15.4	18.3	24	21.9
Cysteine	2.2	2.0	2.2	3.3	3.1
Glutamic acid	45.6	40.6	47	48.5	45
Glycine	16.3	8.4	10.1	12.8	12.6
Serine	13.0	10.4	12.2	14.4	13.2
Tyrosine	8.1	7.6	9.3	10.1	9.5

CPC; corn protein concentrate based diet, CGM; corn gluten meal based diet, HPDDG; high protein distillers dried grain based diet, DDGS; distillers dried grains with soluble based diet; EAA, essential amino acid; NEAA, non essential amino acid

Table 4 Growth performances and feed utilization efficiencies of Nile tilapia fed diets containing corn co-products over 12 weeks (mean $\pm$ SD, n=2)

	Control	CPC	CGM	HPDDG	DDGS	p
% SGR	3.56 $\pm$ 0.01 ^a	2.63 $\pm$ 0.01 ^d	2.75 $\pm$ 0.20 ^c	3.30 $\pm$ 0.03 ^b	3.53 $\pm$ 0.06 ^a	*
TGC	1.21 $\pm$ 0.01 ^a	0.76 $\pm$ 0.01 ^e	0.81 $\pm$ 0.01 ^d	1.06 $\pm$ 0.01 ^c	1.16 $\pm$ 0.00 ^b	*
Feed intake (g dry weight)	84.05 $\pm$ 0.78 ^a	38.80 $\pm$ 0.28 ^b	40.2 $\pm$ 0.14 ^b	71.05 $\pm$ 9.85 ^a	81.20 $\pm$ 1.20 ^a	*
FCR	1.00 $\pm$ 0.00	1.10 $\pm$ 0.00	1.00 $\pm$ 0.00	1.05 $\pm$ 0.15	1.05 $\pm$ 0.07	ns
PER	3.20 $\pm$ 0.01	2.84 $\pm$ 0.11	3.10 $\pm$ 0.08	2.99 $\pm$ 0.50	3.06 $\pm$ 0.06	ns
Protein retention (%)	49.62 $\pm$ 0.13 ^a	38.42 $\pm$ 0.78 ^c	42.02 $\pm$ 1.63 ^{bc}	46.17 $\pm$ 2.91 ^{ab}	46.70 $\pm$ 0.81 ^{ab}	*
% Survival	100.0 $\pm$ 0.0 ^a	75.0 $\pm$ 4.0 ^c	66.6 $\pm$ 7.8 ^c	80.6 $\pm$ 3.9 ^{bc}	97.2 $\pm$ 3.9 ^{ab}	*

Mean values in same row with different superscripts are statistically different (p < 0.05)

% SGR: percentage specific growth rate; TGC : thermal growth coefficient FCR: food conversion ratio; PER: protein efficiency ratio

ns : not significant, *: p < 0.05)

Table 5 Proximate composition of whole body and fillet (% wet basis) of Nile tilapia fed experimental diets over 12 weeks (Mean $\pm$ SD, n=3)

Treatment	Moisture	Protein	Lipid	Ash
<i>Whole body</i>				
Control	69.4 $\pm$ 1.6	15.5 $\pm$ 1.1 ^b	8.5 $\pm$ 0. 6 ^b	6.9 $\pm$ 0. 5 ^a
CPC	71.6 $\pm$ 0.8	13.9 $\pm$ 0.3 ^d	9.6 $\pm$ 1.3 ^a	5.0 $\pm$ 0. 1 ^d
CGM	70.9 $\pm$ 1.0	14. 6 $\pm$ 1.0 ^c	9.8 $\pm$ 2.9 ^a	4.0 $\pm$ 0. 4 ^e
HPDDG	68.9 $\pm$ 0.7	16. 7 $\pm$ 1.9 ^a	9.9 $\pm$ 3.6 ^a	5.4 $\pm$ 1.2 ^c
DDGS	69.7 $\pm$ 1.2	15.4 $\pm$ 0.3 ^b	10.0 $\pm$ 0. 4 ^a	5.7 $\pm$ 0. 4 ^b
<i>Fillet</i>				
Control	78.2 $\pm$ 1.2	18.8 $\pm$ 0. 6 ^b	1.6 $\pm$ 0. 2 ^c	1.4 $\pm$ 0.5
CPC	78.5 $\pm$ 1.6	18.7 $\pm$ 2.8 ^b	1.9 $\pm$ 0. 1 ^{bc}	1.4 $\pm$ 0. 4
CGM	77.9 $\pm$ 0.8	19.2 $\pm$ 2.4 ^b	2.2 $\pm$ 0.8 ^b	1.3 $\pm$ 0.1
HPDDG	76.2 $\pm$ 1.3	19.8 $\pm$ 1.2 ^a	2.4 $\pm$ 0.5 ^b	1.2 $\pm$ 0.7
DDGS	77.2 $\pm$ 0.6	18.3 $\pm$ 0. 9 ^b	3.08 $\pm$ 0.8 ^a	1.3 $\pm$ 0.1

Mean values in same row with different superscripts are statistically different (p < 0.05)

Table 6 Whole body amino acid composition (g kg⁻¹ wet basis) of Nile tilapia fed experimental diets over 12 weeks (Mean $\pm$ SD, n=3).

Amino acid	Control	CPC	CGM	HPDDG	DDGS	p value
EAA						
Arginine	10.3 $\pm$ 1.2	9.1 $\pm$ 0.5	10.2 $\pm$ 2.2	10.8 $\pm$ 0.1	10.2 $\pm$ 1.2	0.747
Histidine	3.4 $\pm$ 0.4	2.9 $\pm$ 0.1	3.4 $\pm$ 0.8	3.4 $\pm$ 0.1	3.4 $\pm$ 0.5	0.758
Isoleucine	5.1 $\pm$ 0.8	4.4 $\pm$ 0.1	5.8 $\pm$ 1.3	4.5 $\pm$ 0.3	5.4 $\pm$ 0.6	0.443
Leucine	11.5 $\pm$ 1.4	10.6 $\pm$ 0.4	12.4 $\pm$ 2.8	11.2 $\pm$ 0.5	11.8 $\pm$ 1.8	0.695
Lysine	10.2 $\pm$ 1.1	9.4 $\pm$ 1.3	10.8 $\pm$ 2.3	11.0 $\pm$ 0.1	10.5 $\pm$ 1.5	0.774
Methionine	3.5 $\pm$ 0.4	3.1 $\pm$ 0.2	4.0 $\pm$ 0.9	3.2 $\pm$ 0.1	3.7 $\pm$ 0.4	0.467
Phenylalanine	6.2 $\pm$ 0.8	5.2 $\pm$ 0.4	6.6 $\pm$ 1.4	5.6 $\pm$ 0.4	6.4 $\pm$ 1.1	0.586
Threonine	7.7 $\pm$ 0.9	6.6 $\pm$ 0.8	8.1 $\pm$ 1.8	7.3 $\pm$ 0.5	7.8 $\pm$ 1.2	0.690
Tryptophan	1.1 $\pm$ 0.2	0.9 $\pm$ 0.1	1.1 $\pm$ 0.2	1.2 $\pm$ 0.4	1.1 $\pm$ 0.4	0.742
Valine	6.2 $\pm$ 0.9	5.4 $\pm$ 0.7	6.8 $\pm$ 1.6	5.7 $\pm$ 1.4	6.4 $\pm$ 0.6	0.537
NEAA						
Alanine	12.2 $\pm$ 1.3	10.5 $\pm$ 0.4	11.8 $\pm$ 2.5	12.8 $\pm$ 0.3	11.9 $\pm$ 1.6	0.642
Aspartic acid	16.2 $\pm$ 1.8	14.2 $\pm$ 0.1	13.4 $\pm$ 2.6	16.4 $\pm$ 0.5	16.3 $\pm$ 2.4	0.405
Cysteine	0.8 $\pm$ 0.1	0.6 $\pm$ 0.2	1.0 $\pm$ 0.1	0.6 $\pm$ 0.1	0.8 $\pm$ 0.2	0.195
Glutamic acid	24.0 $\pm$ 2.6	21.0 $\pm$ 0.1	24.6 $\pm$ 5.3	24.7 $\pm$ 0.4	23.8 $\pm$ 3.3	0.748
Glycine	15.4 $\pm$ 1.9	12.2 $\pm$ 1.1	13.4 $\pm$ 2.9	15.8 $\pm$ 0.6	14.0 $\pm$ 1.3	0.334
Serine	7.7 $\pm$ 0.9	6.6 $\pm$ 0.1	7.9 $\pm$ 1.6	7.6 $\pm$ 0.2	7.8 $\pm$ 1.0	0.688
Taurine	2.1 $\pm$ 0.1	2.8 $\pm$ 0.8	3.0 $\pm$ 0.7	3.8 $\pm$ 0.2	2.7 $\pm$ 0.1	0.151
Tyrosine	5.0 $\pm$ 0.6	4.4 $\pm$ 0.4	5.4 $\pm$ 1.3	4.4 $\pm$ 0.4	5.3 $\pm$ 0.8	0.591

Table 7 Body indices of Nile tilapia fed diets containing corn co-products over 12 weeks ((Mean $\pm$ SD, n=10)

Morphometry index	Control	CPC	CGM	HPDDG	DDGS
VSI	10.8 $\pm$ 0.5	12.8 $\pm$ 1.4	12.1 $\pm$ 0.5	11.6 $\pm$ 0.9	12.9 $\pm$ 1.6
HSI	3 .0 $\pm$ 0.2	2 .0 $\pm$ 0.8	2 .2 $\pm$ 1.4	2.1 $\pm$ 0.1	2.7 $\pm$ 0.7
FY	30.4 $\pm$ 2.6	28.3 $\pm$ 1.1	31.9 $\pm$ 2.6	30.8 $\pm$ 1.1	32.4 $\pm$ 1.9
K	2.0 $\pm$ 0.1	1.9 $\pm$ 0.1	1.8 $\pm$ 0.1	2.0 $\pm$ 0.2	2.0 $\pm$ 0.1

VSI, viscerosomatic index = 100 x visceral weight (g)/ body weight (g); HSI, hepatosomatic index = 100 x liver weight (g)/body weight (g); FY, fillet yield = 100 x fillet weight (g) /body weight (g) ; K, coefficient of condition = 100 x body weight (g)/ total length (cm)³