Functional Sperm of the Yellowtail (*Seriola quinqueradiata*) Were Produced in the Small-Bodied Surrogate, Jack Mackerel (*Trachurus japonicus*).

著者	Morita Tetsuro, Kagayaki Morishima, Misako Miwa, Naoki Kumakura, Satomi Kudo, Ichida Kensuke, Toru Mitsuboshi, Yutaka Takeuchi, Yoshizaki Goro
journal or publication title	Marine Biotechnology
volume	17
number	5
page range	644-654
year	2015-10
権利	(c) 2015 Springer US. This is the author's version of the work. It is posted here for your personal use. Not for redistribution. The definitive Version of Record was published in https://doi.org/10.1007/s10126-015-9657-5
科学研究費研究課題	サケ科魚類の進化に伴うGSC制御機構の変化 Evolution of GSC regulation in salmonids Planned
研究課題番号	25114005
URL	http://id.nii.ac.jp/1342/00001607/

doi: 10.1007/s10126-015-9657-5

Figure legends

Fig. 1 Section of the donor testis of the yellowtail stained with hematoxylin and eosin. **a** Overview of the testis. Upper and lower rectangles show the area rich in type A spermatogonia and sperm, respectively. **b** and **c** show higher magnification images of upper and lower rectangles in **a**, respectively. Scale bars: 2 mm (**a**) and 20 μ m (**b** and **c**).

Fig. 2 Intraperitoneal transplantation of donor testicular cells into the jack mackerel larvae. **a** Bright-field image. **b** Fluorescent image. **c** Transplanted (TP) and non-transplanted (NTP) individuals. N, transplantation needle. Scale bar = 1 mm.

Fig. 3 Incorporation of transplanted PKH26-labeled cells into recipient genital ridges. Fluorescent (**a**, **c**) and yellowtail *vasa in situ* hybridization (**b**, **d**) images of the excised genital ridges from transplanted (**a**, **b**) and nontransplanted (**c**, **d**) fish at 20 dpt. *Arrowheads* indicate donor-derived germ cells that were labeled with PKH-26 (**a**), and were expressing the yellowtail *vasa* mRNA (**b**). Scale bars = 20 μ m.

Fig. 4 Sections of jack mackerel (**a**, **b**) and the yellowtail (**c**, **d**) testis hybridized with antisense jack mackerel (**a**, **c**) and yellowtail *vasa* probes (**b**, **d**). Scale bars = 20 μ m.

Fig. 5 RT-PCR amplification of donor-derived yellowtail *vasa* mRNA in recipient gonads.

Electrophoresis patterns of RT-PCR of recipient number 1–14 (**a**) and 15–28 (**b**) using specific primers for the yellowtail *vasa* (upper panels) and jack mackerel *vasa* (lower panels) cDNA sequences. *Lane P* is a positive control obtained using cDNA derived from the yellowtail testis as template. *Lane N* is a negative control containing no cDNA template. *M* represents a molecular weight marker.

Fig. 6 Detection of the colony containing donor-derived germ cells in recipient testis. Serial sections of testis from jack mackerel testis hybridized with yellowtail (**a**) and jack mackerel (**b**) *vasa* probes and were stained with hematoxylin and eosin (**c**). The colony containing donor-derived germ cells is surrounded with a dotted line. Scale bars = 20 μ m.

Fig. 7 PCR amplification of donor-derived yellowtail *vasa* genomic DNA fragment in recipient semen. *Lane P* is a positive control obtained using genomic DNA extracted from the yellowtail semen as template. *Lane N* is a negative control containing no DNA template. Electrophoretic mobility of the targeted DNA fragment was indicated by *arrowheads*.

Fig. 8 Embryos at 48 h-post-fertilization obtained from progeny tests. **a** Normal embryos obtained by crossing the yellowtail female and male. **b** Eggs obtained by crossing the yellowtail female and male recipient number 28. *Arrowhead* indicates a normal embryo. **c** Eggs obtained by crossing the yellowtail female and wild-type jack mackerel male. Scale bars = 1 mm.

Fig. 9 Donor-derived offspring obtained from progeny tests. **a** Newly hatched larva obtained by crossing the yellowtail female and male. **b–d** Newly hatched larvae obtained by crossing the yellowtail female and male recipient number 28. **b** Normal individual. **c, d** Abnormal individuals. Scale bars = 500 μm .

Fig. 10 PCR amplification of *β -actin* and *vasa* genomic DNA fragments in larvae obtained from progeny tests. Electrophoresis patterns of PCR using specific primers for yellowtail *β -actin* (**a**), jack mackerel *β -actin* (**b**), yellowtail *vasa* (**c**), and jack mackerel *vasa* (**d**) DNA sequences. *Lane 1* and *2* are larva samples obtained from the crosses of recipient mackerel number 28 and the yellowtail females. *Lane 3* and *4* are larva samples of wild type yellowtail. *Lane 5* and *6* are wild type jack mackerel. *Lane 7* is a negative control containing no DNA template. *M* represents a molecular weight marker.