

Isolation of aquatic yeasts with the ability to neutralize acidic media, from an extremely acidic river near Japan's Kusatsu-Shirane Volcano

著者	三矢 太亮, Takuya Hayashi, Yu Wang, Mami Tanaka, 岡井 公彦, 石田 真巳, 浦野 直人
journal or publication title	Journal of Bioscience and Bioengineering
volume	124
number	1
page range	43-46
year	2017-07
権利	(c) 2017, The Society for Biotechnology, Japan. This is the author's version of the work. It is posted here for your personal use. Not for redistribution. The definitive Version of Record was published in https://doi.org/10.1016/j.jbiosc.2017.02.005
科学研究費研究課題	アルカリ化酵母とその酵素系による強酸性水圏中和の分子機構の解明
研究課題番号	16K07868
URL	http://id.nii.ac.jp/1342/00001582/

doi: 10.1016/j.jbiosc.2017.02.005



1 **Table 1.** Identification of the isolates by an ITS1 sequence analysis

Strain	Organism (BLAST result)	Sequence identity ^a	GenBank accession no.
(i) Sampling site upstream of neutralization plant (27.9°C, pH 1.9)			
AeA6	<i>Candida fluviatilis</i>	100% (189/189)	LC133464
EeB28	<i>Candida palmioleophila</i>	100% (191/191)	LC133465
EeC21	<i>Candida fluviatilis</i>	100% (189/189)	LC133466
(ii) Sampling site downstream of neutralization plant (27.8°C, pH 4.9)			
CeA14	<i>Candida fluviatilis</i>	100% (189/189)	LC133467
CeA16	<i>Candida fluviatilis</i>	100% (189/189)	LC133463
CeG17	<i>Candida fluviatilis</i>	100% (189/189)	LC133468
GeC45	<i>Candida fluviatilis</i>	100% (189/189)	LC133469

2 ^a Percentage of identity (Number of identical sequence / aligned sequence shared with
3 type strain). Type strain: *C. fluviatilis* ATCC 38621^T (GenBank accession no.
4 HQ652068, NCBI RefSeq accession no. NR_111589), *C. palmioleophila* CBS 7418^T
5 (GenBank accession no. KY102283, NCBI RefSeq accession no. NR_077076).

6
7