

Functional Sperm of the Yellowtail (*Seriola quinqueradiata*) Were Produced in the Small-Bodied Surrogate, Jack Mackerel (*Trachurus japonicus*).

著者	Tetsuro Morita, Kagayaki Morishima, Misako Miwa, Naoki Kumakura, Satomi Kudo, Ichida Kensuke, Toru Mitsuboshi, Yutaka Takeuchi, Yoshizaki Goro
journal or publication title	Marine Biotechnology
volume	17
number	5
page range	644-654
year	2015-10
権利	(c) 2015 Springer US. This is the author's version of the work. It is posted here for your personal use. Not for redistribution. The definitive Version of Record was published in https://doi.org/10.1007/s10126-015-9657-5
科学研究費研究課題	サケ科魚類の進化に伴うGSC制御機構の変化 Evolution of GSC regulation in salmonids Planned
研究課題番号	25114005
URL	http://id.nii.ac.jp/1342/00001607/

doi: 10.1007/s10126-015-9657-5



1 Table 1. Survival of jack mackerel recipients and colonization of PKH26-positive cells in recipient
 2 genital ridges at 20 days after the xenogeneic transplantation of the yellowtail spermatogonia.

dph at transplantation	Total length (mm)	Group	No. transplanted	No. survived (%)	No. observed ^b	No. colonized ^c
10	3.96	Transplanted	250	134 (53.6)	10	8
		Control ^a	300	238 (79.3)	3	0
12	4.32	Transplanted	273	177 (64.8)	7	7
		Control ^a	300	263 (87.7)	3	0

3 ^a Control means non-transplanted fish group.

4 ^b Number of recipient fish whose genital ridges were observed under fluorescent microscopy.

5 ^c Number of recipient fish whose genital ridges were found to possess the PKH26-labeled cells.

6

7 Table 2. Results of progeny tests and appearance proportions of donor-derived larvae.

Male	Female	No. eggs used for tests	No. eggs fertilized (%) ^c	No. hatchlings (%) ^d	No. donor- derived larvae (%) ^e
Recipient 28	Yellowtail F1	18887	10706 (56.7)	3 (0.028)	1 (0.009)
Recipient 82	Yellowtail F1	21745	9269 (42.6)	3 (0.032)	0 (0.000)
Non transplanted 1 ^a	Yellowtail F1	22700	6965 (30.5)	0 (0.000)	
Yellowtail M1 ^b	Yellowtail F1	5375	4329 (80.5)	3702 (68.9)	
Recipient 28	Yellowtail F2	11385	2980 (26.2)	2 (0.067)	1 (0.033)
Recipient 82	Yellowtail F2	12000	1344 (11.2)	0 (0.000)	0 (0.000)
Non transplanted 2 ^a	Yellowtail F2	11402	4280 (37.5)	0 (0.000)	
Yellowtail M2 ^b	Yellowtail F2	7129	5396 (75.7)	2825 (39.6)	

8 ^aWild type male fish of jack mackerel for non-transplanted control.9 ^bYellowtail male fish for positive control.10 ^cFertilization (%) = number of eggs with cleaved blastodisc/number of eggs used for test × 100.11 ^dHatching (%) = number of newly hatched larvae/number of eggs fertilized × 100.12 ^eGermline transmission (%) = number of donor-derived larvae/number of eggs fertilized × 100.

13

14 Table 3. Genotype pattern of three microsatellite loci in offspring obtained by the progeny tests.

Cross (female × male)	Fish	Microsatellite locus		
		sequ-56	sequ-57	sequ-77
	Donor	94/108	150/161	148/156
	Recipient 28	ND ^a	ND ^a	180/195
Yellowtail F1 × Recipient 28	Yellowtail F1	100/114	125/148	152/156
	Larvae 1	94/100	148/150	148/152
	Larvae 2	100/	148/	152/
	Larvae 3	100/114	125/148	152/156
Yellowtail F2 × Recipient 28	Yellowtail F2	100/	135/	167/210
	Larvae 4	94/100	135/161	156/167
	Larvae 5	100/	135/	167/

15 ^aGenotype were not determined.

16